

Delocalization of two-particle ring near the Fermi level of 2d Anderson model

J. Lages and D.L. Shepelyansky^a

Laboratoire de Physique Quantique^b, Université Paul Sabatier, 31062 Toulouse Cedex 4, France

Received 29 September 2000 and Received in final form 15 January 2001

Abstract. We study analytically and numerically the problem of two particles with a long range attractive interaction on a two-dimensional (2d) lattice with disorder. It is shown that below some critical disorder the interaction creates delocalized coupled states near the Fermi level. These states appear inside well localized noninteracting phase and have a form of two-particle ring which diffusively propagates over the lattice.

PACS. 72.15.Rn Localization effects (Anderson or weak localization) – 71.30.+h Metal-insulator transitions and other electronic transitions – 74.20.-z Theories and models of superconducting state – 05.45.Mt Semiclassical chaos (“quantum chaos”)

Recently a great deal of attention has been attracted to the problem of interaction effects in disordered systems with Anderson localization [1,2]. From the theoretical point of view the problem is rather nontrivial. Indeed, even if a great progress has been reached in the theoretical investigation of the properties of localized eigenstates [3] still the analytical expressions for interaction matrix elements between localized states are lacking. In spite of these theoretical difficulties it has been shown recently that a repulsive or attractive interaction between particles can destroy localization and lead to a propagation of pairs in the noninteracting localized phase. This two interacting particles (TIP) effect has been studied recently by different groups [4–11] and it has been understood that the delocalization of TIP pairs is related to the enhancement of interaction in systems with complex, chaotic eigenstates. Such an enhancement had been already known for parity violation induced by the weak interaction in heavy nuclei [12] where the interaction is typically increased by a factor of thousand. However, since there the two-body interaction is really weak the final result still remains small. On the contrary, for TIP pairs in the localized phase the enhancement of interaction qualitatively changes the dynamics leading to a coherent propagation of TIP on a distance l_c being much larger than the pair size and one-particle localization length l_1 . The enhancement factor κ is determined by the density of two-particle states ρ_2 , coupled by interaction, and the interaction induced transition rate Γ_2 between noninteracting eigenstates, so that $\kappa = \Gamma_2 \rho_2$. At $\kappa \sim 1$ the interaction matrix element

becomes comparable with two-particle level spacing and the Anderson localization starts to be destroyed by interaction. For excited states the TIP density ρ_2 is significantly larger than the one-particle density ρ and the delocalization can be reached for relatively weak interaction if l_1 is large. However, when the excitation energy ϵ above the Fermi level decreases then ρ_2 becomes smaller and it approaches the one-particle density ρ at low energy: $\rho_2 \approx \epsilon \rho^2$ [5]. As a result the value of κ also drops with ϵ so that the delocalization of TIP pairs practically disappears near the Fermi energy. This result has been found in [5,13] in the approximation of the frozen Fermi sea created by fermions. Recent numerical studies of TIP pairs with short range interaction near the Fermi level [14] confirmed these theoretical expectations.

In this paper we discuss another type of situation in which TIP delocalization takes place mainly due to geometrical reasons and not due to the relation $\rho_2 \gg \rho$. As a result the TIP pair can be delocalized in a close vicinity to the Fermi level that opens new interesting possibilities for interaction induced delocalization in the localized noninteracting phase. To study this new situation we investigate a model with a long range attractive interaction between particles in the 2d Anderson model. In this case the particles can rotate around their center of mass, being far from each other and keeping the same energy, while the center can move randomly in two dimensions. As a result the system has effectively three degrees of freedom that makes it rather similar to the case of one particle in the 3d Anderson model where delocalization takes place at sufficiently weak disorder. A somewhat similar situation has been studied recently for particles with Coulomb interaction but only excited states were considered there and the delocalization was attributed to the large ratio

^a e-mail: dima@irsacm2.ups-tlse.fr
<http://w3-phystheo.ups-tlse.fr/~dima>

^b UMR 5626 CNRS

ρ_2/ρ [15]. Here we show that in fact the conditions for delocalization are much less restrictive.

To illustrate the above ideas let us first discuss the case of only two particles with attractive interaction $U(r) < 0$ in the 2d Anderson model described by the Schrödinger equation

$$(E_{\mathbf{n}_1} + E_{\mathbf{n}_2} + U(\mathbf{n}_1 - \mathbf{n}_2))\psi_{\mathbf{n}_1, \mathbf{n}_2} + V(\psi_{\mathbf{n}_1+1, \mathbf{n}_2} + \psi_{\mathbf{n}_1-1, \mathbf{n}_2} + \psi_{\mathbf{n}_1, \mathbf{n}_2+1} + \psi_{\mathbf{n}_1, \mathbf{n}_2-1}) = E\psi_{\mathbf{n}_1, \mathbf{n}_2}. \quad (1)$$

Here $\mathbf{n}_{1,2}$ are the indices of the two particles on the 2d lattice with L^2 sites and periodic boundary conditions, V is the hopping between nearby sites and the random on-site one-particle energies $E_{\mathbf{n}_{1,2}}$ are homogeneously distributed in the interval $[-W/2, W/2]$. The long range attractive interaction depends only on the distance between particles $r = \|\mathbf{n}_1 - \mathbf{n}_2\|$ and is equal to a constant $U < 0$ if $|r - R| \leq \Delta R$ and zero otherwise. The value of r is determined as the minimal inter-particle distance on the periodic lattice. Thus the interaction takes place only inside a ring of radius R and width ΔR , and we assume that $R \gg \Delta R \geq 1$. For $U = 0$ the eigenstates are given by the product of two one-particle (noninteracting) eigenstates which are always localized in 2d in a presence of disorder [16].

In the limit of very strong attractive interaction $|U| \gg V, W$ the TIP coupled states form the energy band of width $\simeq 16V$ around $E \simeq -|U|$ (we consider only the states symmetric in respect to particle interchange). For states in this band the particles are located always inside the ring which center can move over the 2d lattice. Since $|U| \gg V$ these states are decoupled from all other states with particles outside the ring. In the ring the Schrödinger equation is in fact rather similar to the case of 3d Anderson model of one particle. In this analogy the number of sites inside the ring $M_R \approx 2\pi R \Delta R$ determines the effective number of 2d planes placed one over another in the third z -dimension (length size $L_z = M_R$). In this 3d model the effective strength of disorder is approximately $2W$ since the diagonal term is now the sum of two $E_{\mathbf{n}}$ values. Also one site is coupled with $Z = 8$ neighbours contrary to $Z = 6$ for 3d case (assuming $\Delta R \gg 1$). Since in 3d the Anderson transition at the band center takes place at $W_c = 2.75ZV = 16.5V$ [17], we expect that TIP states inside the ring will be delocalized in the middle of the band when $2W/ZV = 2.75$ that gives the transition at $W_{c2} \approx 11V$. The above relation corresponds to the border according to which the mixing of levels takes place when the coupling becomes comparable to the spacing between directly coupled states [13] which in our case is proportional to W/Z with the numerical factor taken from [17]. This estimate is in agreement with preliminary numerical simulations of the model (1) [18]. Of course, since in the slab the thickness in the third direction is finite the eigenstates will be eventually localized. But their localization length l_c will make a sharp jump from $l_c \sim 1$ at $W > W_{c2}$ to $l_c \sim \exp(g) \gg 1$ at $W < W_{c2}$ that follows from the standard scaling theory in 2d [3,16,15]. Here g is the conductance of the quasi-two-dimensional layer of width $L_z = M_R$. As usual $g = E_c/\Delta_1$ where

$E_c = D/L^2$ is the Thouless energy, $\Delta_1 \sim V/(L^2 L_z)$ is the level spacing [3] and the diffusion rate in the lattice model is $D \sim V(V/W)^2$. As a result for $W < W_{c2}$ the TIP delocalization length jumps to exponentially large value $l_c \sim \exp(2\pi R \Delta R (W_{c2}/W)^2)$. In these estimates we assumed that $l_1 > \Delta R > 1$ since if $\Delta R \gg l_1$ the majority of states inside the ring are noninteracting and can be presented as the product of one-particle eigenstates. We also note that for $W < W_{c2}$ there is an energy interval around the band center with delocalized states where the TIP ring diffuses with the rate $D_2 \sim V(W_{c2}/V)^2$. When W decreases the mobility edge approaches the bottom of the band as it happens in 3d Anderson model.

The above arguments presented for the case $|U| \gg V$ indicate that it is possible to have a similar TIP delocalization at moderate value of $U \sim V$ near the Fermi level. To investigate this case we rewrite the equation (1) in the basis of the noninteracting eigenstates that gives

$$(E_{m_1} + E_{m_2})\chi_{m_1, m_2} + U \sum_{m'_1, m'_2} Q_{m_1, m_2, m'_1, m'_2} \chi_{m'_1, m'_2} = E\chi_{m_1, m_2}. \quad (2)$$

Here χ_{m_1, m_2} are eigenfunctions of the TIP problem written in the basis of one-particle eigenstates ϕ_m with eigenenergies E_m . The matrix $UQ_{m_1, m_2, m'_1, m'_2}$ represents the two-body matrix elements of interaction $U(\mathbf{n}_1 - \mathbf{n}_2)$ between noninteracting eigenstates $|\phi_{m_1} \phi_{m_2}\rangle$ and $|\phi_{m'_1} \phi_{m'_2}\rangle$. The Fermi sea is determined by the restriction of the summation in (2) to $m_{1,2}^{(\prime)} > 0$ with energies $E_{m_{1,2}^{(\prime)}} > E_F$, where E_F is the Fermi energy related to the filling factor μ . We choose the case with half filling $\mu = 1/2$ for which $E_F \approx 0$. In this way our model corresponds to the approximation of frozen Fermi sea successfully used for the Cooper problem [19]. As it was done by Cooper we also introduce the high energy cut-off defined by the condition $1 \leq m'_1 + m'_2 \leq M$. This rule determines an effective phonon frequency $\omega_D \propto M/L^2$. We fix $\alpha = L^2/M \approx 15$ since ω_D should be independent of the system size L [20]. We checked that the results are not affected by a variation of α in few times. The first studies of the TIP model with frozen Fermi sea was done by Imry [5] with the aim to take into account the effect of finite fermionic density and then was also analyzed in [13]. The one-dimensional case was also studied in [7]. Recently a similar model was investigated for the case of Hubbard attraction in 3d [14]. Of course, the Cooper approach does not take into account many-body effects. However, it is known that without disorder it captures an important part of physics. Therefore we think that the generalization of this approach to systems with disorder will allow to reach a better understanding of rather complicated many-body problem.

To study the eigenstate properties of our model we diagonalize numerically the Hamiltonian (2) and rewrite the eigenfunctions in the original lattice basis. In this way we determine the two-particle probability distribution $F(\mathbf{n}_1, \mathbf{n}_2)$ from which we extract the one particle




Fig. 1. Probability distributions f and f_d for TIP in 2d disordered lattice of size $L = 40$, and interaction of radius $R = 12$ and width $\Delta R = 1$. Left column, one-particle probability f for $W = 8V$: (a) ground state at $U = 0$; (b) ground state with binding energy $\Delta E = -1.05V$ at $U = -2V$; (c) coupled state with $\Delta E = -0.19V$ at $U = -2V$. Right column: (d) f for coupled state, compare to case (c), at $W = 12V$ and $U = -2V$ with $\Delta E \approx -0.19V$; (e) inter-particle distance probability f_d related to case (b); (f) f_d related to case (c).

probability $f(\mathbf{n}_1) = \sum_{\mathbf{n}_2} F(\mathbf{n}_1, \mathbf{n}_2)$ and the probability of inter-particle distance $f_d(\mathbf{r}) = \sum_{\mathbf{n}_2} F(\mathbf{r} + \mathbf{n}_2, \mathbf{n}_2)$ with

$\mathbf{r} = \mathbf{n}_1 - \mathbf{n}_2$. The binding energy of an eigenstate in (2) is $\Delta E = E - 2E_F \approx E$ since $E_F \approx 0$. For the ground state with energy E_g the coupling energy is $\Delta = 2E_F - E_g$. The typical examples of probability distributions are shown in Figure 1. They clearly show that the ground state in the presence of interaction remains localized (b) and the particles stay on distance R from each other (e). However, there are states with negative binding energy ($\Delta E < 0$) which are delocalized by interaction (c) and for which the particles move around the ring (f) in agreement with discussion of model (1) at $|U| \gg V$. We stress that this delocalization of coupled states ($\Delta E < 0$) takes place in the well localized one-particle phase. However, at very strong disorder this delocalization disappears (see top right case in Fig. 1d).

To analyze the delocalization of states with negative binding energy ΔE in a more quantitative way we determine the inverse participating ratio (IPR) ξ for one-particle probability $1/\xi = \left\langle \sum_{\mathbf{n}} f^2(\mathbf{n}) \right\rangle$, where




Fig. 2. 2d density plots for the data of Figure 1 with the same ordering of figures (color figures are available at <http://xxx.lanl.gov/abs/cond-mat/0002296>). Black corresponds to the minimum of the probability distribution and light-grey to the maximum. Figures (a, b, d, e) are drawn in logarithmic scale while Figures (c, f) are in linear scale. Black/light-grey color corresponds to: (a) $f = 1.3 \times 10^{-10}/f = 0.2$, (b) $f = 1.1 \times 10^{-9}/f = 0.26$, (c) $f = 1.5 \times 10^{-6}/f = 0.014$, (d) $f = 1.14 \times 10^{-11}/f = 0.073$, (e) $f_d = 3.8 \times 10^{-7}/f_d = 0.1$, (f) $f_d = 1.5 \times 10^{-5}/f_d = 0.0032$.

brackets mark the averaging over 100 disorder realizations. In this way ξ gives the number of lattice sites occupied by one particle in an eigenstate. The dependence of ξ on ΔE and W is shown in Figure 3 for different lattice sizes L in the presence of interaction. This figure shows that near the ground state the interaction creates states which are even more localized than in the absence of interaction (ξ is significantly smaller than at $U = 0$, see insert Fig. 3). In fact for $-\Delta < \Delta E < \Delta E_c < 0$ the IPR value even slightly drops with the increase of L . However for the states with binding energy $\Delta E_c < \Delta E < 0$ the situation becomes different and ξ grows significantly with L while the change of IPR at $U = 0$ with L is rather weak (see shaded band in Fig. 3). The critical value of the binding energy ΔE_c can be defined as such an energy at which ξ remains independent of L . In this way ΔE_c determines




Fig. 3. Dependence of IPR ξ on the binding energy $\Delta E/V$ for $U = -2V$, $W = 8V$ and $\Delta R = 1$: $L = 20$ (\circ), $L = 22$ (triangle down), $L = 26$ (diamond), $L = 30$ (triangle up), $L = 40$ (square); full/empty symbols are for $R = 8/R = 12$; the shaded band shows the variation of ξ at E_F and $U = 0$ for $20 \leq L \leq 40$. The delocalization transition at ΔE_c is marked by arrows. Insert shows dependence of ξ on disorder W/V at $R = 8$ for the ground state at $U = -2V$, $L = 30$ (\times) and at $U = 0$, $L = 30$ ($+$); for the states with binding energy ΔE_c at $U = -2V$ the ξ value is shown by ($*$).

the mobility edge for coupled states so that at given U and W the TIP eigenstates are localized for $-\Delta < \Delta E < \Delta E_c$ while for $\Delta E_c < \Delta E < 0$ the states becomes delocalized (see an example in Fig. 1). In agreement with this picture ξ varies up to 30 times when ΔE changes from $-\Delta$ up to 0. This variation grows with L and the interaction radius R since the system becomes more close to the effective 3d Anderson model as it was discussed above. The qualitative change of the structure of the eigenstates leads also to a change in the level spacing statistics $P(s)$ (Fig. 4). Near the ground state the statistics is close to the Poisson distribution $P_P(s) = \exp(-s)$ typical for the localized Anderson phase [21] while for $\Delta E_c < \Delta E < 0$ it approaches to the Wigner surmise $P_W(s) = \pi s \exp(-\pi s^2/4)/2$ corresponding to the delocalized phase [21].

The variation of the delocalization border ΔE_c for TIP coupled states with disorder strength and interaction is shown in Figure 5. While the coupling energy Δ grows with U and W , the mobility edge $\Delta E_c < 0$, on the contrary, disappears at strong W . According to the data of Figure 5 all states with binding energy $\Delta E < 0$ become localized for $W > W_{c2} \approx 9.5V$ ($U = -2V$) and $W > W_{c2} \approx 8V$ ($U = -V$). This shows that at weaker interaction a weaker disorder is required to have delocalized coupled states. We note that the values of W_{c2} obtained for $U = -V, -2V$ are rather close to the value $W_{c2} \approx 11$ corresponding to the limit $|U| \gg V, W$. This shows that the physics of delocalization at $U \sim V$ remains rather similar to the limiting case of very strong interaction. As it follows from Figure 5, at small disorder W the delocalization border ΔE_c becomes closer and closer to the ground




Fig. 4. Level spacing statistics $P(s)$ for TIP coupled states at $L = 30$, $U = -2V$, $W = 8V$, $R = 8$, $\Delta R = 1$ in the localized phase near the ground state inside the energy interval $-\Delta < \Delta E < -3\Delta/4$ (\circ) and in the delocalized phase for energies inside $-\Delta/4 < \Delta E < 0$ (\bullet); here $\Delta E_c \approx -0.3\Delta$ and the statistics is done over 3000 disorder realizations with the total number of spacings being 3×10^4 (\circ)/ 7.5×10^4 (\bullet). Full lines show the Poisson distribution and the Wigner surmise.




Fig. 5. Dependence of the delocalization border ΔE_c and the binding energy of the ground state $-\Delta$ on disorder W/V . The values of ΔE_c are shown by (\circ)/(\bullet) for $U = -V / -2V$; the values of $-\Delta$ are shown by upper/lower dashed line for $U = -V / -2V$ and $L = 30$ (vertical intervals give variation for $20 \leq L \leq 30$).

state. This means that at weak disorder the delocalization will take place for excited states with low energy. For $W \ll W_{c2}$ and $U \sim -V$ the diffusion rate of delocalized TIP ring can be estimated as $D_2 \sim V(W_{c2}/W)^2$ [22]. Further studies are required to determine the dependence of W_{c2} on W at $|U| \ll V$.

In conclusion, our results show that long range attractive interaction between two particles in 2d leads to the appearance of delocalized diffusive states near the Fermi level inside the well localized noninteracting phase. It would be interesting to understand what will be the consequences of this delocalization for real many-body fermionic problem with attractive interaction. It is possible that obtained results will be also relevant for electrons with Coulomb repulsion. Indeed, in this case at very

weak disorder each electron oscillates near an equilibrium position and the two-body interaction can be considered as an effective harmonic attraction [23]. Another physical situation which can be related to the investigated model corresponds to the case of long range attractive Coulomb interaction between electron-hole excitation in a semiconductor (*e.g.* spin-exciton). In a strong magnetic field electron/hole rotates on a cyclotron circle which drifts slowly along another large circle formed by an equipotential line of Coulomb interaction. Thus without disorder such a system corresponds very closely to the ring model we analyzed above with the ring width proportional to the cyclotron radius. In the presence of disorder the displacements of the ring can be localized by quantum effects or become delocalized below some critical disorder strength. While this physical situation still should be studied in more detail it is clear that its physical properties are closely related to the ring model analyzed in the paper.

We thank O.P. Sushkov for stimulating discussions, and the IDRIS in Orsay and the CICT in Toulouse for access to their supercomputers.

References

1. *Correlated fermions and transport in mesoscopic systems*, edited by T. Martin, G. Montambaux, J. Trân Thanh Vân (Editions Frontières, Gif-sur-Yvette, 1996).
2. *Disorder and Interaction in Transport Phenomena*, edited by M. Schreiber, Ann. Phys. (Leipzig) **8**, 539–798 (1999).
3. A. Mirlin, Phys. Rep. **328**, 259 (2000).
4. D.L. Shepelyansky, Phys. Rev. Lett. **73**, 2607 (1994); p. 201 in [1].
5. Y. Imry, Europhys. Lett. **30**, 405 (1995); p. 211 in [1].
6. D. Weinmann, A. Müller-Groeling, J.-L. Pichard, K. Frahm, Phys. Rev. Lett. **75**, 1598 (1995); p. 221 in [1].
7. F. von Oppen, T. Wettig, J. Müller, Phys. Rev. Lett. **76**, 491 (1996); p. 235 in [1].
8. P.H. Song, D. Kim, Phys. Rev. B **56**, 12217 (1997).
9. K.M. Frahm, Eur. Phys. J. B **10**, 371 (1998).
10. M. Ortuño, E. Cuevas, Europhys. Lett. **46**, 224 (1999); E. Cuevas, Phys. Rev. Lett. **83** (1999) 140.
11. R.A. Römer, M. Leadbeater, M. Schreiber, p. 675 in [2].
12. O.P. Sushkov, V.V. Flambaum, Usp. Fiz. Nauk. **136**, 3 (1982) [Sov. Phys. Usp. **25**, 1 (1982)].
13. Ph. Jacquod, D.L. Shepelyansky, Phys. Rev. Lett. **78**, 4986 (1997).
14. J. Lages, D.L. Shepelyansky, Phys. Rev. B **62**, 8665 (2000).
15. D.L. Shepelyansky, Phys. Rev. B **61**, 4588 (2000).
16. E. Abrahams, P.W. Anderson, D.C. Licciardello, T.V. Ramakrishnan, Phys. Rev. Lett. **42** 673 (1979).
17. T. Ohtsuki, K. Slevin, T. Kawarabayashi, p. 655 in [2].
18. G. Caldara, D.L. Shepelyansky (unpublished).
19. L.N. Cooper, Phys. Rev. **104**, 1189 (1956).
20. In this case we found that $\omega_D/V \approx 0.055W/V + 0.20$.
21. B.I. Shklovskii, B. Shapiro, B.R. Sears, P. Lambrianides, H.B. Shore, Phys. Rev. B **47**, 11487 (1993).
22. We expect that D_2 will drop with ΔE growth from ΔE_c to 0; indeed, for $\Delta E > 0$ the particles are in average more separated from each other and their interaction becomes effectively smaller.
23. V.V. Flambaum, I.V. Ponomarev, O.P. Sushkov, Phys. Rev. B **59**, 4163 (1999).